

# Adagolási útmutató



LATIN	ENGLISH	PROCONT	REVITA-U	ERGERON	REVITASOL	ADDITIONAL PRODUCT	COMMENTS
Dosage given: (time * number of tablets)			(female 2*1, from week 2. 3*1) (male 2*1)	*2, in case of bad values 3 (Consume the pro- duct in the evening!)	**1, in case of bad values 2		
In the table, the increased dosing is marked in red!							
Carcinoma cervicis uteri, (HPV)	Cervical Cancer	3*2	3*1	2	2		
Carcinoma laryngis	Laryngeal Cancer	3*2	2*1	2	2		
Carcinoma mammae	Breast Cancer	„2-1-2; 3*2”	2*1	2	2		Procont dos.: if hormone sensitive 2-1-2, if triple negat. 3*2
Carcinoma pancreatis	Pancreatic tumor	3*2	2*1	3	2		In most cases, metastasis can occur in the liver!
Carcinoma ventriculi	Stomach tumor	3*2	3*1	3	2		Probably liver metastasis, very rapid course! Immediate treatment's required!
Chirrhosis Hepatis	Cirrhosis	3*1	3*1	4	2		
Colorectalis carcinoma (CRC)	Colon cancer, Rectal cancer	3*2	3*1	„2-3”	2		In most cases there's liver metastasis, ERGERON dosage depending on it!
Glioblastoma	Brain Tumor	3*2	2*1	2	1		
Hepatocellularis carcinoma (Hepatitis B, C) Liver Cancer		„2-2-3”	3*1	„3-4”	2		
Melanoma	Melanoma	3*2	2*1	2	2		
Mioma, Endometriosis	Mioma	2*2	3*1	2	1	2*2 AQUARES	Use an additional product!
Prostata carcinoma	Prostatic Cancer	„2*2;3*2” Based on PSA value	3*1	2	1	2*1;3*1 PROSTANOVA 7 (Based on PSA value) Zinc cone	Frequent pelvic metastases. Use an additional product!
Pulmo adeno carcinoma	Lung tumor	3*2	2*1	2	2	2*1 AEROVITAL	Use an additional product!
Pulmo carcinoma microcellulare	Lung small cell tumor	„2-2-3”	3*1	2	2	2*1 AEROVITAL	Use an additional product!
Renal carcinoma	Kidney tumor	3*2	2*1	2	1	2*1 RENALONUM	Use an additional product!
Thiyeoidea carcinoma	Thyroid tumor	3*2	2*1	2	2	I*1 THYR. CONT. Should be taken after 1 hour of hormone pills!	Use an additional product!
Multiplex metastasis	Multiplex metastasis	3*2	3*1	3	2		
Ossealis metastasis	Bone metastasis	3*2	3*1	2	2	3*1 REVITACART	Use an additional product!

\* Blood count based on liver enzyme values (GOT, GPT, GGT, AP, LDH)

\*\* Based on blood count (abs. Lymphocyte, abs. Monocyte,% lymphocyte, red blood cell count, white blood cell count, hemoglobin)

For all tumor types in case of need for support of hematopoietic organs (Based on red blood cell level, Hemoglobin level): 1\*1; 2\*1 REVITAFERR

For all types of tumors to relieve anxiety symptoms: 1-3\*1-2 SEDALON

For all tumor types as a support for restful sleep: 1\*1-2 SOMNIFERON

# Adagolási útmutató



ENGLISH	LATIN	PROCONT	REVITA-U (female-2*1, from week 2- 3*1) (male-2*1)	ERGERON <b>*2, in case of bad values 3 (Consume the pro- duct in the evening!)</b>	REVITASOL <b>**1, in case of bad values 2</b>	ADDITIONAL PRODUCT	COMMENTS
Dosage given: (time*number of tablets)							
Type 2 diabetes	Diabetes 2		2*1		1	„3*1;2*2” AQUARES	Aquares dosage depends on blood sugar, (over 10 – 2*2)
Stroke	Stroke		2*1		1	2*2 AQUARES	Aquares dosage: in the 1th week 2*1, from the 2nd week 2*2
Depression	Depression		2*1		1	3*1;4*1 SEDALON	
Gastric ulcer, Duodenal ulcer	Ulcus ventriculi, Duodenal ulcer	2*1	2*1; 2*2	1	1		
Chronic arthritis	Rheumatoid arthritis (RA, PCP)		2*1; 3*1		1	3*1 REVITACART	
Chronic colitis Chron's disease	Colitis ulcerosa, Chron	3*1	2*1; 3*1; 2*2	1	2		
Hypertension	Hypertonia			2	1	„3*1;2*2” AQUARES	Aquares dosage is blood pressure dependent, over 160/100 - 2*2
High blood pressure (coronary spasm)	Angina pectoris			1	1	„3*1;2*2” AQUARES	Aquares dosage is blood pressure dependent, over 160/100 - 2*3
Thyroid disorder	Hypothyreosis; Hyperthyreosis					I*1 THYROID CONTROL	Take the Supplement 1 hour after the hormone tablets!
Parkinson's disease	Parkinson	2*1	2*1	2	2	2*2 AQUARES	Elevated Revitasol values!
Sclerosis Multiplex (SM)	Sclerosis multiplex (SM)	2*1	2*1	2	2	2*2 AQUARES	Elevated Revitasol values!
Heart attack	Myocardialis infarctus		2*1		1	2*2 AQUARES	Aquares dosage: in the 1th week 2*1, from the 2nd week 2*2

